


Worksheet 2-1: Probability Experiments

Probability is a measure of the likelihood that a specific event will occur.

Probability experiments can be used to estimate the probability of an event.

Let's use a probability problem to learn about specific terms related to probability.

Problem: A spinner has 4 equal sectors colored yellow, blue, green and red.
 What are the chances of landing on blue after spinning the spinner?
 What are the chances of landing on red?



Solution: The chances of landing on blue are 1 in 4, or one fourth.
 The chances of landing on red are 1 in 4, or one fourth.

Let's look at some definitions and examples from the problem above.

Definition	Example
An experiment is a situation involving chance or probability that leads to results called outcomes.	In the problem above, the experiment is spinning the spinner.
An outcome is a possible result of a single trial of an experiment.	The possible outcomes are landing on yellow, blue, green or red.
An event is one or more outcomes of an experiment.	One event of this experiment is landing on blue.
A trial is one round of a probability experiment	One trial of this experiment is one spin.
Probability is the measure of how likely an event is.	The probability of landing on blue is one fourth.

In order to measure probabilities, mathematicians have devised the following formulas for finding the probability and experimental probability of an event.

Probability Of An Event
$P(\text{Event A}) = \frac{\text{Number of Ways Event A Can Occur}}{\text{Total Number of Possible Outcomes}}$


Experimental Probability of An Event
$P(\text{Event A}) = \frac{\text{Number of Successful Trials}}{\text{Total Number of Trials}}$

Probability is always a value between 0 (*certain not to happen*) and 1 (*certain to happen*).

Probability can be expressed as a fraction in lowest terms, a decimal or a percent.

Assigned Work: WS 2-1; p. 66 #1-4

1. The results of rolling a six-sided die are displayed in the graph.




- (a) How many times was a 5 rolled?
- (b) Find the experimental probability of rolling a 6. Express your answer as a fraction in lowest terms, as a decimal, and as a percent.
- (c) Find the experimental probability of not rolling a 6. How is this related to the probability of rolling a 6?

2. What is the probability of rolling a 7 with a six-sided die?

3. What is the probability of rolling a 1, 2, 3, 4, 5, or 6?

4. A probability experiment was designed to find the expected number of females in a family of six children. To simulate the genders of the six children, a coin was tossed six times. Heads represented a male; tails represented a female. The experiment was repeated a number of times. The results are shown in the graph.



- (a) How many trials were performed?

- (b) What is the experimental probability of having two females in a family of six children?

5. Sam tossed a coin 10 times and heads turned up nine times.

(a) Express the experimental probability of turning up heads as a percent.

(b) Is the experimental probability what you expected? Explain.

6. A coin was tossed 20 times. The experimental probability of turning up tails was 30%.

(a) How many times did the coin turn up tails?

(b) How many times did the coin turn up heads?

(c) What was the experimental probability of the coin turning up heads? Describe two different methods of finding the answer.

Answers: 1. (a) 3 times, (b) $\frac{1}{5}$, 0.2, 20%, (c) $\frac{4}{5}$; 2. 0; 3. 1; 4. (a) 17, (b) $\frac{4}{17}$; 5. (a) 90%, (b) No, expect 50% because the two outcomes are equally likely; 6. (a) 6 times, (b) 14 times, (c) 70% ($100\% - 30\%$ or $\frac{14}{20} \times 100$).